

The Role of DNA Damage and Extracellular Trap Formation in Atherosclerosis

Vickie Tang

*A thesis submitted in fulfilment of the requirements for the award
of the degree Doctor of Philosophy (Science)*

October, 2020

School of Maths and Physical Sciences

Faculty of Science

University of Technology Sydney

Certificate of Original Authorship

I, Vickie Tang declare that this thesis, is submitted in fulfillment of the requirements for the award of PhD: Science, in the School of Maths and Physical Sciences at the University of Technology Sydney.

This thesis is wholly my own work unless otherwise referenced or acknowledged. In addition, I certify that all information sources and literature used are indicated in the thesis.

This thesis contains materials that have been published in [1] and is shown in Chapter 3. This publication includes the results shown in Figures 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 3.10 and 3.11. I designed all the studies, analysed the data and wrote the drafts for the manuscript.

This document has not been submitted for qualifications at any other academic institutions.

This research is supported by the Australian Government Research Training Program.

Vickie Tang

Production Note:
Signature removed prior to publication.

31st October 2020

*"Nothing in life is to be feared, it is only to be understood.
Now is the time to understand more, so that we may fear less."*

- **Marie Curie**

"Even miracles take a little time."

- **Fairy Godmother**

*"You have brains in your head. You have feet in your shoes.
You can steer yourself any direction you choose.
You're on your own. And you know what you know.
And you are the one who'll decide where to go..."*

- **Dr Seuss**

Acknowledgements

Firstly, a huge thank you must go to my supervisors Prof Clare Hawkins, Prof Shanlin Fu and Dr Benjamin Rayner for providing constant support throughout my PhD. They have truly gone above and beyond to provide experience, guidance and a wealth of knowledge throughout this journey. Similarly, thanks must also go to the other members of the Inflammation Group at the Heart Research Institute and the Redox Biology Group at the University of Copenhagen. The members of both these groups provided insight, advice, personal experience and most importantly an empathetic shoulder to cry on when everything seemed to be going wrong.

Thanks must also go to people who were more than willing to share their experimental expertise with me. To Dr Allan Weimann and Prof Henrik Poulsen, thank you for sharing your knowledge in the detection of oxidized nucleosides with me. Also, thanks to Dr Dominic Love for teaching me how to perform the Seahorse experiments. A special thanks must also go to Pat Pisansarakit for all the tissue culture help over the years.

Throughout my PhD candidature, I was lucky enough to be surrounded by the most wonderful friends. Without their unwavering support, I would not have escaped with what's left of my sanity. To my friends I met in my time at the University of Copenhagen, thank you for making my move to a new country so seamless and memorable. My gratitude goes to the crew at the University of Technology Sydney, who have provided nothing but laughter and support. More specifically, my biggest thanks must go to Zohra,

who I was lucky enough to meet during our Honours year. Since then, she has been there for every lunch break and more importantly as emotional support. To my best friend, Shani, we have survived another massive milestone in our lives, and I couldn't be more thankful to have had you by my side as we've lived our best life together for the past 10 years. To my male twin and fellow gin lover, Ali, I will always be so grateful to have met you during the most stressful period of our lives. You have provided much needed encouragement and for that, I'll forever be indebted to you.

My most heartfelt thanks go to my family for whom I will be eternally grateful. To my parents, Selina and Thomas, thank you for providing nothing but love and support throughout the years. To Carrie and David, I will be ever so thankful for your company after a long day and for your constant push to finish this thesis (not to mention, thanks for also providing the best distractions, Joshua and Charlotte). Last but definitely not least, to my now husband, Justin, without whom this thesis truly would not exist. I am forever grateful for the company that you have provided, your poor attempts at distracting me from the mounting stress and for listening to me vent my frustrations even when you had no idea what I was talking about.

Table of Contents

Certificate of Original Authorship.....	i
Acknowledgements	iii
Table of Contents	v
Abstract.....	xii
List of Figures	xiv
List of Tables.....	xvii
Abbreviations	xviii
Publications and conference presentations arising from this thesis	xxiv
Publications	xxiv
Poster presentations	xxv
Oral presentations.....	xxvi
Awards received	xxvii
Chapter 1 – Introduction.....	1
1.1. Overview.....	1
1.2. Cardiovascular disease and atherosclerosis.....	2
1.2.1. Significance of cardiovascular disease	2
1.2.2. Mechanisms responsible for the development and progression of atherosclerosis.....	3
1.3. Cellular dysfunction involved in atherosclerosis.....	6
1.3.1. Endothelial dysfunction	6
1.3.2. Smooth muscle cell dysfunction	8
1.3.3. Macrophages and foam cell formation.....	9

1.3.4.	Neutrophils and neutrophil extracellular trap formation.....	11
1.4.	Myeloperoxidase and atherosclerosis	15
1.4.1.	Chlorinated oxidant production.....	15
1.4.2.	MPO in atherosclerosis	16
1.4.3.	HOCl-induced modifications of proteins.....	19
1.4.4.	Reactions of hypochlorous acid with lipids.....	20
1.4.5.	HOCl-induced modifications of nucleic acids.....	22
1.5.	Biomarkers of inflammation in disease.....	24
1.5.1.	DNA damage products as biomarkers	24
1.5.2.	Detection of DNA damage products	26
1.6.	DNA damage in disease	30
1.6.1.	Cancer	30
1.6.2.	Diabetes	31
1.6.3.	CVD and atherosclerosis	32
1.7.	Biological reactivity of chlorinated nucleosides.....	33
1.7.1.	Cellular consequences.....	33
1.7.2.	Chlorinated nucleosides as therapeutic agents.....	35
1.8.	Summary.....	37
1.9.	Hypotheses and aims	38
1.9.1.	Hypotheses.....	38
1.9.2.	Project aims.....	39
Chapter 2 – Materials and Methods		40
2.1.	Materials.....	40
2.2.	Chlorinated nucleosides and all-trans retinoic acid (ATRA).....	40
2.3.	Cell culture.....	40

2.3.1.	Human coronary artery endothelial cell culture.....	40
2.3.2.	Human coronary artery smooth muscle cell culture	41
2.3.3.	PLB-985 cell culture and differentiation	42
2.3.3.1.	Determination of PLB-985 morphology with May-Grünwald-Giemsa staining.....	42
2.4.	Neutrophil extracellular trap release	43
2.4.1.	Qualitative detection of NETs by microscopy with Sytox Green staining	43
2.4.2.	Quantification of NETs by fluorescence.....	44
2.4.2.1.	Quantification of NETs by Sytox Green staining	44
2.4.2.2.	Quantification of NETs by PicoGreen staining	45
2.5.	Quantification of unmodified and modified nucleosides using liquid chromatography triple quadrupole mass spectrometry (LC-MS/MS)	46
2.5.1.	Nucleic acid isolation	46
2.5.1.1.	Isolation of DNA, RNA and protein from HCAEC and HCASMC	46
2.5.1.2.	Isolation of DNA from extracellular traps released by PLB-985	47
2.5.2.	Hydrolysis and dephosphorylation of DNA and RNA.....	48
2.5.3.	Detection of chlorinated and parent nucleosides by LC-MS/MS.....	49
2.5.4.	Detection of 8CldG in NETs.....	51
2.6.	Quantification of unmodified and chlorinated nucleotides using high performance liquid chromatography (HPLC)	52
2.6.1.	Lysis of HCAEC and HCASMC.....	52
2.6.2.	Extraction of the nucleotide pool	52
2.6.3.	Detection of chlorinated nucleotides	53
2.6.4.	Standard curves and quantification of parent and chlorinated nucleotides	53
2.7.	Cellular assays.....	54

2.7.1.	Metabolic activity by PrestoBlue Cell Viability Reagent	54
2.7.2.	Proliferation assay.....	54
2.7.3.	ATP assay.....	55
2.7.4.	BCA assay	55
2.7.5.	GAPDH activity assay.....	56
2.7.6.	Glucose uptake assay	56
2.7.7.	Seahorse assay	57
2.7.8.	DC protein assay.....	59
2.7.9.	Lactate assay	59
2.7.10.	MMP activity assay	60
2.8.	Flow cytometry.....	61
2.8.1.	Fluo-4AM – Calcium accumulation in the cytosol	61
2.8.2.	Annexin V/propidium iodide – Cell death by apoptosis and necrosis	62
2.8.3.	KI-67 – Cell proliferation	63
2.8.3.1.	Cell treatment and ethanol fixation.....	63
2.8.3.2.	Sample staining with KI-67.....	63
2.8.4.	JC-1 – Mitochondrial membrane depolarisation	64
2.9.	Gene expression by quantitative real time polymerase chain reaction	65
2.9.1.	Cell lysis and extraction of RNA	65
2.9.2.	RNA quantification	66
2.9.3.	cDNA synthesis.....	67
2.9.4.	Quantitative real time polymerase chain reaction (qPCR)	67
2.10.	Protein expression	69
2.10.1.	Lysis of whole cell samples	69
2.10.2.	Nuclear and cytoplasmic extraction.....	69

2.10.3. Reducing protein samples and gel electrophoresis	70
2.10.4. Protein probing, imaging and quantification	71
2.11. Statistical analysis	73
Chapter 3 – Effect of chlorinated nucleosides on endothelial cell function	74
3.1. Introduction.....	74
3.2. Hypothesis and aims.....	77
3.3. Methods	77
3.4. Results	80
3.4.1. Chlorinated nucleosides incorporate into the cellular DNA and RNA	80
3.4.2. 8CIA does not alter ATP production by oxidative phosphorylation in HCAEC	83
3.4.3. Exposure to 8CIA causes decreases ATP synthesis by glycolysis	87
3.4.4. 8CIA treatment does not alter glucose uptake in HCAEC but decreases GAPDH activity	88
3.4.5. Treatment with 8CIA causes ER stress	91
3.4.6. Exposure to 8CIA alters antioxidant responses in HCAEC.....	94
3.4.7. 8CIA causes increased calcium accumulation and induces apoptotic cell death	98
3.5. Discussion	101
3.6. Summary.....	107
Chapter 4 – Effect of chlorinated nucleosides on human coronary artery smooth muscle cells	109
4.1. Introduction.....	109
4.2. Hypothesis and aims.....	112
4.3. Methods	112
4.4. Results	114

4.4.1.	Chlorinated nucleosides incorporate into HCASMC DNA and RNA	114
4.4.2.	Metabolic activity of HCASMC decreases with exposure to 8ClA, 8ClG, 8ClIdG and 5ClIdC	117
4.4.3.	Exposure to 8ClA and 5ClIdC decreases proliferation of HCASMC	119
4.4.4.	8ClA and 5ClIdC do not cause apoptosis or necrosis in HCASMC	122
4.4.5.	8ClA and 5ClIdC do not cause calcium accumulation in HCASMC	123
4.4.6.	8ClA decreases mitochondrial function in HCASMC	124
4.4.7.	8ClA causes the decrease in cellular ATP through its incorporation to form 8-ClATP	125
4.4.8.	Exposure to 8ClA causes activation of the UPR and antioxidant responses in HCASMC	128
4.4.9.	5ClIdC caused increased expression of MMP2 and MMP9 in HCASMC ..	131
4.4.10.	HCASMC exposed to 5ClIdC have increased expression of adhesion molecules	136
4.4.11.	Exposure of HCASMC to 5ClIdC induces phenotypic changes	138
4.5.	Discussion	141
4.6.	Summary	150
Chapter 5 – DNA damage in neutrophil extracellular trap formation		151
5.1.	Introduction	151
5.2.	Hypothesis and aims	154
5.3.	Methods	155
5.4.	Results	156
5.4.1.	Differentiation of PLB-985 cells into neutrophils	156
5.4.2.	Optimisation of PMA-induced and HOCl-induced NET production in differentiated PLB-985	158

5.4.3.	Quantification of NETs produced by PLB-985 using Sytox Green reagent	160
5.4.4.	Quantification of NETs produced by PLB-985 using PicoGreen reagent	162
5.4.5.	Detection and quantification of parent and modified deoxyribose nucleosides in NETs and cellular DNA	163
5.5.	Discussion	168
5.6.	Summary	174
Chapter 6 – General Discussion		176
6.1.	Overview	176
6.2.	Sources and detection of modified DNA and RNA	177
6.3.	Biological impacts of chlorinated nucleosides on cell signalling and viability	183
6.4.	Future directions	190
6.5.	Final conclusions	191
References		194
Appendix 1 – Table of Materials		232
Appendix 2 – Example chromatograms from LC-MS/MS		239

Abstract

The primary cause of cardiovascular disease is atherosclerosis, which is an inflammatory disease resulting in plaque formation within blood vessels. Myeloperoxidase (MPO), an enzyme released by activated leukocytes in atherosclerosis catalyses the formation of hypochlorous acid (HOCl) which reacts readily with nucleic acids forming chlorinated nucleosides. There is increasing evidence that the formation of these chlorinated adducts has adverse effects, and elevated levels of these species are present in different inflammatory contexts, including atherosclerotic lesions.

The focus of Chapters 3 and 4 in this Thesis was to examine the effects of chlorinated nucleosides on endothelial cells (HCAEC) and smooth muscle cells (HCASMC) isolated from the human coronary artery. Studies in HCAEC and HCASMC showed that each of the chlorinated nucleosides studied incorporated into the RNA and DNA of these cells. However, only 8-chloroadenosine (8ClA) was shown to influence the metabolic activity and cell viability in HCAEC. 8ClA was converted to 8-ClATP, which was associated with alterations in the ability of HCAEC to perform glycolysis. Exposure of HCAEC to 8ClA also induced endoplasmic reticulum (ER) stress, resulting in apoptotic cell death and activation of the Nrf2 antioxidant stress response.

In Chapter 4, studies in HCASMC demonstrated that 8ClA and 5-chloro-2'-deoxycytidine (5ClIdC) decreased both the metabolic activity and proliferation of HCASMC. The effects of 8ClA on HCASMC were similar to those observed in HCAEC though less profound as the viability of HCASMC was unaffected, though some ER stress was apparent. In contrast, exposure of HCASMC to 5ClIdC resulted in alterations to pro-inflammatory

signalling, a large increase in the expression of MMP2 and MMP9 and phenotypic changes.

A potential source of chlorinated nucleosides under inflammatory conditions was explored in Chapter 5. The differentiation of PLB-985 cells into neutrophils was optimised, together with various treatment conditions to induce the production of neutrophil extracellular traps (NETs). The concentrations of modified nucleosides in both NET DNA and nuclear DNA of these cells was quantified using LC-MS/MS and it was shown that phorbol myristate acetate (PMA) and HOCl treatment gave 8-oxo-7,8-dihydro-deoxyguanosine (8oxodG) and 5ClidC, respectively, in both NET and nuclear DNA.

Overall, the studies outlined in this Thesis provide insight into a potential new source of modified nucleosides in disease. These data show that chlorinated nucleosides are not only biomarkers of disease but can alter cellular function in both HCAEC and HCASMC. Therefore, these chlorinated adducts could play a role in the development and progression of atherosclerosis.

List of Figures

Figure 1.1: Overview of the key steps involved in the development and progression of a plaque in atherosclerosis, reproduced from [39].	6
Figure 1.2: Suicidal and vital neutrophil extracellular trap formation, reproduced from [74].	13
Figure 1.3: Schematic of the formation of HOCl from myeloperoxidase.	15
Figure 1.4: Structures of chlorinated nucleosides.	24
Figure 2.1: Measures of mitochondrial respiration using a Seahorse Analyser (modified from [273]).	58
Figure 2.2: Representative population and quadrant gating for necrosis and apoptosis.	63
Figure 2.3: Representative population gating for JC-1 staining to determine mitochondrial membrane depolarisation.	65
Figure 3.1: Chlorinated nucleosides incorporate into the DNA and RNA of HCAEC.	82
Figure 3.2: The effect of 8CIA on the mitochondrial respiration of HCAEC.	84
Figure 3.3: Mitochondrial respiration remains unaltered after treatment with 8CIA.	86
Figure 3.4: Treatment of HCAEC with 8CIA induces changes in glycolysis and GAPDH activity.	88
Figure 3.5: The effect on glucose uptake when HCAEC are exposed to 8CIA.	90
Figure 3.6: The effect 8CIA on GAPDH activity in HCAEC.	91
Figure 3.7: Treatment with 8CIA causes ER stress and activates the unfolded protein response.	93
Figure 3.8: HCAEC exposed to 8CIA have altered expression of Nrf2-related and other antioxidant responses.	95
Figure 3.9: Treatment with 8CIA does not alter some antioxidant responses in HCAEC.	97
Figure 3.10: Exposure of HCAEC to 8CIA causes calcium to accumulate in the cytosol.	99
Figure 3.11: 8CIA exposure induces apoptotic cell death in HCAEC.	100

Figure 4.1: Accumulation and clearance of chlorinated nucleosides into the DNA and RNA of HCASMC.	116
Figure 4.2: Alterations in metabolic activity on HCASMC exposed to chlorinated nucleosides.....	118
Figure 4.3: Proliferation of HCASMC when exposed to chlorinated nucleosides.	120
Figure 4.4: Expression of KI-67 in HCASMC exposed to chlorinated and parent nucleoside.	121
Figure 4.5: Effects of 8ClA and 5Cl dC on cell death by apoptosis and necrosis.	123
Figure 4.6: Effects of 8ClA and 5Cl dC on calcium accumulation in the cytosol of HCASMC.	124
Figure 4.7: Effects of 8ClA and 5Cl dC on mitochondrial membrane depolarisation....	125
Figure 4.8: Treatment with 8ClA causes the formation of ATP analogue, 8Cl-ATP.....	127
Figure 4.9: Alterations on mRNA expression of genes related to the unfolded protein response.	129
Figure 4.10: The mRNA expression of genes relating to antioxidant response.	130
Figure 4.11: mRNA expression of MMP2 and MMP9 in HCASMC exposed to 8ClA and Ad.	132
Figure 4.12: mRNA expression of MMP2 and MMP9 in HCASMC exposed to 5Cl dC and dC.	134
Figure 4.13: Proteinase activity of MMP2 and MMP9 in HCASMC exposed to 8ClA and 5Cl dC.	135
Figure 4.14: Expression of adhesion molecules after HCASMC were exposed to 5Cl dC and dC.	137
Figure 4.15: mRNA expression of synthetic smooth muscle cell phenotypic markers.	139
Figure 4.16: mRNA expression of contractile smooth muscle cell phenotypic markers.	140
Figure 5.1: May-Grünwald-Giemsa staining of differentiated PLB-985.	157
Figure 5.2: Sytox Green staining of PMA and HOCl-induced extracellular trap release.	159
Figure 5.3: Quantification of NET production by Sytox Green plate reader assay.....	161

Figure 5.4: Quantification of NET production in PLB-985 by PicoGreen plate reader assay.	162
Figure 5.5: Quantification of deoxyribose nucleosides from PMA-induced NETs and nuclear DNA.	165
Figure 5.6: Quantification of deoxyribose nucleosides from HOCl-induced NETs and nuclear DNA.	167

List of Tables

Table 1.1: Mass fragmentation of chlorinated nucleosides detected using LC-MS.	27
Table 2.1: Human coronary artery endothelial cell donor information.	41
Table 2.2: Human coronary artery smooth muscle cell donor information.	42
Table 2.3: Mobile phase composition to separate chlorinated and parent compounds.	50
Table 2.4: Mass fragmentation of unmodified and chlorinated nucleosides.	50
Table 2.5: Gradient elution of 8CldG.	51
Table 2.6: qPCR forward and reverse primer sequences.	68

Abbreviations

Abbreviation	Definition
18S	18S ribosomal RNA
2-NBDG	2-(N-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)amino)-2-deoxyglucose
3-ClTyr	3-chlorotyrosine
5ClC	5-chlorocytidine
5Cl dC	5-chloro-2'-deoxycytidine
5Cl dU	5-chloro-2'-deoxyuridine
5ClU	5-chlorouridine
5ClUra	5-chlorouracil
8ClA	8-chloroadenosine
8-ClATP	8-chloro-ATP
8Cl-cAMP	8-chloro-cyclic AMP
8Cl dA	8-chloro-2'-deoxyadenosine
8Cl dG	8-chloro-2'-deoxyguanosine
8ClG	8-chloroguanosine
8oxodG	8-oxo-7,8-dihydro-deoxyguanosine
8oxoG	8-oxo-7,8-dihydroguanosine
Ad	Adenosine
AMP	Adenosine monophosphate
ANOVA	Analysis of variance
ApoB-100	Apolipoprotein B-100

ApoE ^{-/-}	Apolipoprotein E knockout mice
APS	Ammonium persulfate
ATF4	Activating transcription factor 4
ATF6	Activating transcription factor 6
ATP	Adenosine triphosphate
ATRA	All-trans retinoic acid
BCA	Bicinchoninic acid
BSA	Bovine serum albumin
CCCP	Carbonyl cyanide m-chlorophenyl hydrazone
cDNA	Complimentary DNA
CHOP	CCAAT-enhancer-binding protein homologous protein
CTP	Cytidine triphosphate
CVD	Cardiovascular disease
CX43	Connexin 43
Cyt	Cytidine
dC	2'-deoxycytidine
dG	2'-deoxyguanosine
DMEM	Dulbecco's Modified Eagle Medium
DMF	Dimethylformamide
DMSO	Dimethyl sulfoxide
DNA	Deoxyribonucleic acid
DNase	Deoxyribonuclease
ECAR	Extracellular acidification rate

ECD	Electrochemical detection
EDTA	Ethylenediaminetetraacetic acid
ELISA	Enzyme-linked immunosorbent assay
eNOS	Endothelial nitric oxide synthase
ER	Endoplasmic reticulum
ESI	Electrospray ionisation
ETC	Electron transport chain
FBS	Fetal bovine serum
FCCP	Carbonyl cyanide-4-(trifluoromethoxy)phenylhydrazone
FITC	Fluorescein isothiocyanate
FPG	Formamidopyrimidine DNA glycosylase
GADD34	Growth arrest and DNA damage-inducible protein 34
GAPDH	Glyceraldehyde 3-phosphate dehydrogenase
GCLC	Glutamate-cysteine ligase catalytic subunit
GC-MS	Gas chromatography mass spectrometry
GPx4	Glutathione peroxidase 4
GS	Glutathione synthetase
GTP	Guanosine triphosphate
HBSS	Hank's balanced salt solution
HCAEC	Human coronary artery endothelial cells
HCASMC	Human coronary artery smooth muscle cells
HDL	High-density lipoprotein
HEPES	4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid

HO-1	Heme oxygenase 1
HPLC	High performance liquid chromatography
ICAM-1	Intercellular adhesion molecule-1
IFN γ	Interferon γ
IL-1	Interleukin-1
IL-10	Interleukin-10
IL-12	Interleukin-12
IL-13	Interleukin-13
IL-1 β	Interleukin-1 β
IL-4	Interleukin-4
IL-6	Interleukin-6
IL-8	Interleukin-8
IRE1	Inositol-requiring enzyme 1
Keap-1	Kelch-like ECH-associated protein 1
LC-MS	Liquid chromatography mass spectrometry
LC-MS/MS	Triple quadrupole liquid chromatography mass spectrometry
LDL	Low density lipoprotein
LOD	Limit of detection
LPS	Lipopolysaccharide
M1	Pro-inflammatory macrophage
M2	Anti-inflammatory macrophage
MMP2	Matrix metalloproteinase 2
MMP9	Matrix metalloproteinase 9

Mnase	Micrococcal nuclease
MPO	Myeloperoxidase
MRM	Multiple reaction monitoring
mRNA	Messenger RNA
NADH	Nicotinamide adenine dinucleotide
NADPH	Nicotinamide adenine dinucleotide phosphate
NETs	Neutrophil extracellular traps
NMR	Nuclear magnetic resonance
NQO1	NADPH quinone dehydrogenase 1
Nrf2	Nuclear factor erythroid-2-related factor 2
OCR	Oxygen consumption rate
Olfml3	Olfactomedin Like 3
OPN	Osteopontin
oxLDL	Oxidised low-density lipoprotein
PAD4	Peptidylarginine deiminase 4
PBS	Phosphate buffered saline
PERK	PKR-like endoplasmic reticulum kinase
PI	Propidium iodide
PMA	Phorbol 12-myristate 13-acetate
PVDF	Polyvinylidene fluoride
qPCR	Quantitative real time polymerase chain reaction
RIPA	Radioimmunoprecipitation assay
RNA	Ribonucleic acid

ROS	Reactive oxygen species
RPL13	Ribosomal protein L13
RPMI	Roswell Park Memorial Institute medium
S100A4	S100 calcium binding protein A4
SDS	Sodium dodecyl sulphate
SEM	Standard error of the mean
SOD1	Superoxide dismutase 1
SOD2	Superoxide dismutase 2
SR-AI	Scavenger receptor-AI
SR-AII	Scavenger receptor-AII
sXBP1	Spliced X-box binding protein 1
TBST	Tris-buffered saline with 0.1% Tween 20
TEMED	Tetramethylethylenediamine
TFA	Trifluoroacetic acid
TLR	Toll-like receptor
TNF α	Tumour necrosis factor α
TRAIL	Tumour necrosis factor related apoptosis inducing ligand
Trx	Thioredoxin
TXNIP	Thioredoxin interacting protein
UDG	Uracil DNA glycosylase
UPR	Unfolded protein response
UTP	Uridine triphosphate
VCAM-1	Vascular cell adhesion molecule-1

Publications and conference presentations arising from this thesis

Publications

Tang, V., Macer-Wright, J., Fu, S., Rayner, B., Hawkins, C., *Chlorinated Nucleosides - A Novel Inducer of Endothelial Dysfunction in Atherosclerosis?* Free Radical Biology and Medicine, 2016. **100**: p. S152.

Tang, V., Macer-Wright, J., Fu, S., Rayner, B., Hawkins, C., *A role for chlorinated nucleosides in the promotion of inflammation and endothelial dysfunction in atherosclerosis?* Free Radical Biology and Medicine, 2017. **108**: p. S30.

Tang, V., Macer-Wright, J., Rayner, B., Fu, S., Hawkins, C., *The effect of chlorinated nucleosides on vascular cell function.* Free Radical Biology and Medicine, 2018. **120**: p. S92-S93.

Tang, V., Rayner, B., Fu, S., Hawkins, C., *Role of chlorinated nucleosides in the promotion of vascular smooth muscle cell dysfunction.* Free Radical Biology and Medicine, 2018. **128**: p. S103-S104.

Tang, V., Fu, S., Rayner, B., Hawkins, C., *8-Chloroadenosine induces apoptosis in human coronary artery endothelial cells through the activation of the unfolded protein response.* Redox Biology, 2019. **26**: p. 101274.

Poster presentations

Tang, V., Macer-Wright, J., Fu, S., Rayner, B., Hawkins, C.

“Chlorinated Nucleosides - A Novel Inducer of Endothelial Dysfunction in Atherosclerosis?”

23rd Annual Meeting of Society for Redox Biology and Medicine, 16th – 19th November 2016, San Francisco, USA.

Tang, V., Macer-Wright, J., Fu, S., Rayner, B., Hawkins, C.

“A role for chlorinated nucleosides in the promotion of inflammation and endothelial dysfunction in atherosclerosis?”

OCC World Congress and Annual Society for Free Radical Research Europe, 21st – 23rd June 2017, Berlin, Germany.

Tang, V., Macer-Wright, J., Rayner, B., Fu, S., Hawkins, C.

“The effect of chlorinated nucleosides on vascular cell function.”

19th Biennial Meeting of Society for Free Radical Research International, 4th – 7th June 2018, Lisbon, Portugal.

Tang, V., Rayner, B., Fu, S., Hawkins, C.

“Role of chlorinated nucleosides in the promotion of vascular smooth muscle cell dysfunction.”

25th Annual Meeting of Society for Redox Biology and Medicine, 14th – 17th November 2018, Chicago, USA

Oral presentations

Tang, V., Macer-Wright, J., Fu, S., Rayner, B., Hawkins, C.

“The role of chlorinated nucleosides in endothelial dysfunction and atherosclerosis.”

**2017 Australian Society for Medical Research NSW Annual Scientific Meeting, 2nd June
2017, Sydney, Australia.**

Tang, V., Macer-Wright, J., Fu, S., Rayner, B., Hawkins, C.

*“A role for chlorinated nucleosides in the development of endothelial dysfunction in
atherosclerosis?”*

**2017 Australian Atherosclerosis Society Annual Scientific Meeting, 25th – 27th October
2017, Sydney, Australia.**

Tang, V., Rayner, B., Fu, S., Hawkins, C.

*“Chlorinated nucleosides and their role in endothelial and smooth muscle cell
dysfunction.”*

**8th Joint Meeting of Society for Free Radical Research Australasia and Japan, 9th – 12th
December 2017, Tokyo, Japan.**

Awards received

Student Oral Finalist, Australian Atherosclerosis Society 2017, Sydney, Australia.

Young Investigator Finalist, Australian Society for Medical Research 2017, Sydney, Australia.

Young Investigator Travel Award, Society for Free Radical Research Australasia 2017, Tokyo, Japan.

Young Investigator Travel Award, Society for Free Radical Research Australasia 2018, Lisbon, Portugal.

July Paper of the Month, University of Technology Sydney 2019, Sydney, Australia.